

Non-Abelian Antisymmetric-Vector Coupling from Self-Interaction

Adel Khoudeir¹

*Centro de Astrofísica Teórica, Departamento de Física, Facultad de
Ciencias, Universidad de los Andes, Mérida, 5101, Venezuela.*

Abstract

A non-abelian coupling between antisymmetric fields and Yang-Mills fields proposed by Freedman and Townsend several years ago is derived using the self-interaction mechanism.

1 INTRODUCTION

Abelian second-rank antisymmetric fields [1] play an essential role in strings and supergravity theories and have been extensively studied in the last decades [2] [3] [4] [5]. In free theories they describe massless and spinless particles and appear in many contexts, for instance, arising as mediators of the interaction between open strings with charged particles [2] and in ten dimensions, coupling with the Chern-Simons 3-form to achieve an elegant unification of Yang-Mills and supergravity [6]. In particular the Cremmer-Sherk theory [3] has received considerable attention [7] [8] due to the fact that the coupling between the abelian antisymmetric field and a Maxwellian field through a topological BF term leads to massive propagations which are compatible with gauge invariances. Moreover, Allen, et. al. [7] have shown unitarity and renormalizability of the Cremmer-Sherk theory. This fact motivates the non-abelian generalization of the model and several attempts have been proposed [9]. Simultaneously, other alternatives for non-abelian massive vector bosons without the presence of Higgs field have been proposed in the last year [10].

The non-abelian extension of antisymmetric theories was achieved by Freedman and Townsend [4] starting from a first-order formulation where

¹ e-mail: adel@ciens.ula.ve

the antisymmetric field B_{mn}^a and an auxiliary vector potential are independent variables. It is worth recalling that the non-abelian generalization of the abelian S-duality theory [11] is a Freedman- Townsend theory [12]. In their work, Freedman and Townsend proposed the non-abelian generalization of the Cremmer-Sherk theory. In this letter, starting from an appropriate first-order formulation for the Cremmer-Sherk theory, we will derive the non-abelian generalization using the self-interaction mechanism [13], which has been successfully applied to formulate Yang-Mills, gravity [13], supergravity [14], topologically massive Yang-Mills [15] and Chapline-Manton [16] theories.

2 THE ABELIAN MODEL

Our starting point will be a first-order formulation for the Cremmer-Sherk theory. This is realized introducing an auxiliary vector field (v_m) *ala* Freedman-Townsend. The action is written down as [17]

$$I = < -\frac{1}{4}\mu\epsilon^{mnpq}B_{mn}[\partial_p v_q - \partial_q v_p] - \frac{1}{2}\mu^2 v^m v_m - \frac{1}{2}\mu\epsilon^{mnpq}B_{mn}\partial_p A_q \quad (1) \\ + \frac{1}{4}F_{mn}F^{mn} - \frac{1}{2}F^{mn}[\partial_m A_n - \partial_n A_m] >$$

where $<>$ denotes integration in four dimensions. All the fields involved have mass dimensions and μ is a mass parameter. There are two sets of abelian gauge invariances:

$$\delta_\lambda A_m = \partial_m \lambda, \quad \delta_\lambda F_{mn} = 0 \quad (2)$$

$$\delta_\zeta B_{mn} = \partial_m \zeta_n - \partial_n \zeta_m, \quad \delta_\zeta v_m = 0. \quad (3)$$

Independent variations in v_m, B_{mn}, F_{mn} and A_m lead to the following equations of motion

$$v^m = -\frac{1}{6\mu}\epsilon^{mnpq}H_{npq}, \quad (4)$$

$$\epsilon^{mnpq}\partial_p[v_q + A_q] = 0, \quad (5)$$

$$F_{mn} = \partial_m A_n - \partial_n A_m, \quad (6)$$

$$\partial_p F^{pm} = \frac{1}{6}\mu\epsilon^{mnpq}H_{npq} \quad (7)$$

where $H_{mnp} \equiv \partial_m B_{np} + \partial_n B_{pm} + \partial_p B_{mn}$ is the field strength associated with the antisymmetric field. The Cremmer-Sherk action is obtained after substituting equations (4) and (6) in (1):

$$I_{CrSc} = -\frac{1}{4}F_{mn[A]}F_{[A]}^{mn} - \frac{1}{12}H_{mnp[B]}H_{[B]}^{mnp} - \frac{1}{4}\mu\epsilon^{mnpq}B_{mn}F_{pq[A]}. \quad (8)$$

On the other hand, equation (5) can be solved (locally) for the v field,

$$v_m = -[A_m + \frac{1}{\mu}\partial_m\phi], \quad (9)$$

where ϕ is a scalar field. Substituting this solution in the action I , the Stuckelberg formulation for massive abelian vector bosons is obtained

$$I_{St} = -\frac{1}{4}F_{mn[A]}F_{[A]}^{mn} - \frac{1}{2}\mu^2[A_m + \frac{1}{\mu}\partial_m\phi][A^m + \frac{1}{\mu}\partial^m\phi]. \quad (10)$$

As it is well known, both formulations (Stuckelberg and Cremmer-Sherk) are equivalent descriptions of massive abelian gauge invariant vectorial theories and propagate three degrees of freedom. This equivalence is reflected by the fact that they are connected by duality [18]. Indeed, since the scalar field appears in equation (10) only through its derivative, we can apply the dualization method due to Nicolai and Townsend [19], which consist in replacing $\partial_m\phi$ by $\frac{1}{2}l_m$ and adding a new term to equation (10): $\epsilon B\partial l$, i.e.

$$I_{Stmod} = -\frac{1}{4}F_{mn[A]}F_{[A]}^{mn} - \frac{1}{2}\mu^2[A_m + \frac{1}{2\mu}l_m][A^m + \frac{1}{2\mu}l_m] + \frac{1}{4}\epsilon^{mnpq}B_{mn}\partial_p l_q. \quad (11)$$

At this stage, B_{mn} is a Lagrange multiplier forcing the constraint $\partial_m l_n - \partial_n l_m = 0$ whose local solution is $l_m = 2\partial_m\phi$. Now, if we eliminate l_m via its equation of motion

$$l^m = \frac{1}{3}\epsilon^{mnpq}H_{npq} - 2\mu A^m \quad (12)$$

and go back to equation (11), the Cremmer-Sherk action is recovered.

Finally, let us recall that the second-order field equations can be written as

$$\partial_p F^{pm} = J^m, \quad \partial_p H^{pmn} = J^{mn}, \quad (13)$$

where

$$J^m = \frac{1}{6}\mu\epsilon^{mnpq}H_{npq} \quad \text{and} \quad J^{mn} = \frac{1}{2}\mu\epsilon^{mnpq}F_{pq} \quad (14)$$

are "topological" currents in the sense that they are conserved without using the equations of motion.

3 THE SELF-INTERACTION PROCESS

Now, we extend the first-order action, equation (1), by introducing a triplet of free abelian antisymmetric fields B_{mn}^a coupled with a triplet of free abelian vector fields A_m^a , ($a = 1, 2, 3$)

$$I_o = < -\frac{1}{4}\mu\epsilon^{mnpq}B_{mn}^a[\partial_p v_q^a - \partial_q v_p^a] - \frac{1}{2}\mu^2 v^{am}v_m^a - \frac{1}{2}\mu\epsilon^{mnpq}B_{mn}^a\partial_p A_q^a \quad (15) \\ + \frac{1}{4}F_{mn}^a F^{amn} - \frac{1}{2}F^{amn}[\partial_m A_n^a - \partial_n A_m^a] >$$

Besides the local gauge transformations

$$\delta_\lambda A_m^a = \partial_m \lambda^a, \quad \delta_\lambda F_{mn}^a = 0 \quad (16)$$

$$\delta_\zeta B_{mn}^a = \partial_m \zeta_n^a - \partial_n \zeta_m^a, \quad \delta_\zeta v_m^a = 0, \quad (17)$$

our action has two global invariances: one is a global $SU(2)$ rotation and the other is a global symmetry associated with the Freedman-Townsend theory:

$$(I) \quad \delta_\omega X^a = g_1 \epsilon^{abc} X^b \omega^c \quad (18)$$

where $X^a = (A_m^a, F_{mn}^a, v_m^a, B_{mn}^a)$ and

$$(II) \quad \delta_\rho B_{mn}^a = g_2 \epsilon^{abc} [v_m^b + A_m^b] \rho_n^c - m \leftrightarrow n, \quad (19) \\ \delta_\rho v_m^a = \delta_\rho A_m^a = \delta_\rho F_{mn}^a = 0,$$

ω and ρ being global parameters. In principle the coupling constants g_1 and g_2 are different. We note that under type II transformations the action changes by a total derivative. The Noether currents associated to these invariances are given by

$$g_1^{-1} j^{am} = \epsilon^{abc} F^{bmn} A_n^c + \frac{1}{2} \mu \epsilon^{mnpq} \epsilon^{abc} B_{pq}^b [A_n^c + v_n^c] \quad (20)$$

and

$$g_2^{-1} K^{amn} = \frac{1}{2} \mu \epsilon^{mnpq} \epsilon^{abc} [A_p^b + v_p^b] [A_q^c + v_q^c]. \quad (21)$$

These are conserved on-shell. In order to couple these currents to the action I_o we must add the corresponding self-interaction terms: I_1 and I_2 defined by:

$$j^{am} \equiv \frac{\delta I_1}{\delta A_m^a}; \quad K^{amn} \equiv -2 \frac{\delta I_2}{\delta B_{mn}^a}. \quad (22)$$

These functional differential equations can easily be integrated. In fact, we find that

$$\begin{aligned} I_1 &= -g_1 < \frac{1}{2}\epsilon^{abc}F^{amn}A_m^bA_n^c + \frac{1}{4}\mu\epsilon^{mnpq}\epsilon^{abc}B_{mn}^aA_p^bA_q^c \\ &+ \frac{1}{2}\mu\epsilon^{mnpq}\epsilon^{abc}B_{mn}^aA_p^bv_q^c > \end{aligned} \quad (23)$$

and

$$\begin{aligned} I_2 &= -g_2 < \frac{1}{4}\epsilon^{mnpq}\epsilon^{abc}B^{amn}v_p^bv_q^c + \frac{1}{4}\mu\epsilon^{mnpq}\epsilon^{abc}B_{mn}^aA_p^bA_q^c \\ &+ \frac{1}{2}\mu\epsilon^{mnpq}\epsilon^{abc}B_{mn}^aA_p^bv_q^c > \end{aligned} \quad (24)$$

However, these two terms have overlapping parts. This situation is akin to what happens in the derivation of supergravity from self-interaction [14]. In order to overcome this obstacle we must require equality of the coupling constants: $g \equiv g_1 = g_2$ and write down the self-interaction action as

$$\begin{aligned} I_{SI} &\equiv -g < \frac{1}{2}\epsilon^{abc}F^{amn}A_m^bA_n^c + \frac{1}{4}\epsilon^{mnpq}\epsilon^{abc}B_{mn}^av_p^bv_q^c \\ &+ \frac{1}{4}\mu\epsilon^{mnpq}\epsilon^{abc}B_{mn}^aA_p^bA_q^c + \frac{1}{2}\mu\epsilon^{mnpq}\epsilon^{abc}B_{mn}^aA_p^bv_q^c > \end{aligned} \quad (25)$$

Actually, we have that

$$j^{am} \equiv \frac{\delta I_{SI}}{\delta A_m^a} \quad \text{and} \quad K^{amn} \equiv -2 \frac{\delta I_{SI}}{\delta B_{mn}^a}. \quad (26)$$

The self-interaction mechanism stops here since no other derivative terms appear in I_{SI} . Finally, the full non-abelian theory is

$$\begin{aligned} I &= I_o + I_{SI} \\ &= < -\frac{1}{4}\mu\epsilon^{mnpq}B_{mn}^a[F_{pq}^a + f_{pq}^a + 2\epsilon^{abc}A_p^bv_q^c] - \frac{1}{2}\mu^2v_m^av^am - \frac{1}{4}F_{mn}^aF^{amn} >, \end{aligned} \quad (27)$$

where

$$F_{mn}^a \equiv \partial_m A_n^a - \partial_n A_m^a + g\epsilon^{abc}A_m^bA_n^c \quad (28)$$

and

$$f_{mn}^a \equiv \partial_m v_n^a - \partial_n v_m^a + g\epsilon^{abc} v_m^b v_n^c \quad (29)$$

which is just that proposed by Freedman and Townsend (equation (2.15) in their paper). As usual, the self-interaction process combines the abelian gauge transformations with the global ones giving rise to non-abelian local gauge transformations. In our case, we have

$$\delta_\alpha A_m^a = \partial_m \alpha^a + g\epsilon^{abc} A_m^b \alpha^c \quad (30)$$

$$\delta_\alpha B_{mn}^a = g\epsilon^{abc} B_{mn}^b \alpha^c \quad (31)$$

$$\delta_\alpha v_m^a = g\epsilon^{abc} v_m^b \alpha^c$$

and

$$\begin{aligned} \delta_\xi B_{mm}^a &= \partial_m \xi^a + g\epsilon^{abc} [A_m^b + v_m^b] \xi^c - m \leftrightarrow n \\ \delta_\xi A_m^a &= 0 = \delta_\xi v_m^a. \end{aligned} \quad (32)$$

The action of Freedman-Townsend, equation (27), is equivalent to massive Yang-Mills (locally) as can be shown after elimination of B_{mn}^a through its equation of motion, which said us that $A_m + v_m$ is a pure gauge.

4 CONCLUSION

In this letter, by starting with a nice abelian first-order formulation, and through the application of the self-interaction mechanism we have obtained the Freedman-Townsend theory and its corresponding gauge tranformation rules through self-interaction. The first order abelian formulation allowed us to find Cremmer-Sherk and Stuckelberg formulations for massive spin-1 theories, these later formulations are connected by duality. The BRST quantization of the massive Freedman-Townsend has been performed by Thierry-Mieg [20]. Since massive Freedman-Townsend theory is equivalent (in topologically trivial manifolds) to massive Yang Mills it should be interesting to attempt to connect Friedman-Townsend with others approaches dealing with massive gauge bosons without the presence of Higgs field [10].

5 ACKNOWLEDGEMENT

I thank P.J. Arias for useful discussions and U. Percoco, M. Caicedo, N. Pantoja and L. Labrador also for carefully reading the manuscript.

6 REFERENCES

References

- [1] V.I. Ogievetsky and I.V. Polubarinov, Sov. J. Nucl. Phys. **4** (1967) 156; S. Deser, Phys. Rev. **187** (1969) 1931.
- [2] M. Kalb and P. Ramond, Phys. Rev. **D9** (1974) 2273.
- [3] E. Cremmer and J. Scherk, Nucl. Phys. **bf B72** (1974) 117.
- [4] D.Z. Freedman and P.K. Townsend, Nucl. Phys. **bf B177** (1981) 282.
- [5] S. Deser and E. Witten, Nucl. Phys. **B178** (1981) 491; S. Deser, P.K. Townsend and W. Siegel, Nucl. Phys. **B184** (1981) 333; J. Thierry-Mieg and L. Baulieu, Nucl. Phys. **B228** (1983) 259.
- [6] E. Bergshoeff, M. de Roo, B. de Wit and P. van Nieuwenhuizen, Nucl. Phys. **B195** (1982) 97; G. F. Chapline and N.S. Manton, Phys. Lett. **B120** (1983) 105.
- [7] T. J. Allen, M. J. Bowick and A. Lahiri, Mod. Phys. Lett. **A6** (1991) 559.
- [8] A. Lahiri, Mod. Phys. Lett. **A8** (1993) 2403; R. Amorin and J. Barcelos-Neto, Mod. Phys. Lett. **A10** (1995) 917.
- [9] A. Lahiri, hep-th/9301060; D. S. Hwang, C.Y. Lee, hep-th/ 9512216; A. Lahiri, hep-th/9602009; J. Barcelos-Neto, A. Cabo and M.B.D. Silva, preprint IC/95/245.
- [10] V. Periwal, hep-th/9509084 and hep-th/950985; J. de Boer, K. Skenderis and P. van Nieuwenhuizen, Phys. Lett. **B367** (1996) 175; A. J. Niemi, Phys. Lett. **B374** (1996) 119.

- [11] E. Witten, hep-th/9506011
- [12] Y. Lozano, Phys. Lett. **B364** (1995) 19; N. Mohammedi hep-th/9507040; C. Hong-Mo, J. Faridani and T.S. Tsun, hep-th/9503106; O. Ganor and J. Sonnenschein, hep-th/9507036
- [13] S. Deser, Gen. Rel. Grav. **1** (1970) 9; Class. Quant. Grav. **4** (1987) L99.
- [14] D. G. Boulware, S. Deser and J.H. Kay Physics **96A** (1979) 141.
- [15] C. Aragone and E. Araujo, Acta Científica Venezolana, **36** (1985) 207.
- [16] C. Aragone and J. Stephany, Rev. Bras. Fis. **16** (1987) 287; Quantum Mechanics of Fundamental Systems 1, edited by C. Teitelboim(Plenum Press, N.Y, 1988) p. 27.
- [17] We use the metric $\eta_{mn} = diag(-1, +1, +1, +1)$.
- [18] E. Harikumar and M. Sivakumar, hep-th/9604181.
- [19] H. Nicolai and P.K. Townsend, Phys. Lett. **B98** (1981) 257.
- [20] J. Thierry-Mieg, Nucl. Phys. **B335** (1990) 334.